



Product Datasheet

Product Name	Recombinant Human Retinol Binding Protein-4, His tag
Cata No	CB501289
Source	<i>Escherichia Coli</i> .
Synonyms	Retinol Binding Protein 4 plasma, RBP-4, RBP4, Plasma retinol-binding protein, PRBP, RBP.

Description

Retinol binding protein 4(RBP4) belongs to the lipocalin family and is the specific carrier for retinol (vitamin A alcohol) in the blood. This protein was found to be expressed and secreted by adipose tissue, and was strongly associated with insulin resistance. It delivers retinol from the liver stores to the peripheral tissues. In plasma, the RBP-retinol complex interacts with transthyretin which prevents its loss by filtration through the kidney glomeruli. RBP4 delivers retinol from the liver to the peripheral tissues. In plasma, the rbp-retinol complex interacts with transthyretin, this prevents its loss by filtration through the kidney glomeruli.

RBP-4 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 204 amino acids and having a molecular mass of 23 kDa.

RBP4 is fused to N-terminus His tag which contains additional 21 amino acids. RBP4 sequence is identical to UniProtKB/Swiss-Prot entry Q5VY30 amino acids 16–199.

The Retinol Binding Protein-4 is purified by proprietary chromatographic techniques.

Physical Appearance

White lyophilized (freeze-dried) powder.

Biological Activity

Determined by its ability to bind all trans retinol

acids. The binding of RBP4 to all trans retinol acids results in quenching of Trp fluorescence. 0.5M retinol acids/per M RBP4.

Purity

Greater than 95.0% as determined by:

- (a) Analysis by RP-HPLC.
- (b) Analysis by SDS-PAGE.

Formulation

RBP-4 His tag was filtered (0.4 µm) and lyophilized from 0.5 mg/ml in PBS buffer, pH 7.5.

Reconstitution

It is recommended to reconstitute the lyophilized Retinol binding protein-4 in sterile deionized water to a working volume of 0.5mg/mL and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.

Stability

Lyophilized Retinol binding protein-4 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Retinol binding protein-4 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

Please prevent freeze-thaw cycles.

*** For Non-Clinical Research Use Only ***



California Bioscience

83103 Avenue 48, Ste.1B #204
Coachella, CA 92236 USA
Phone : +1.6268339877
Email : info@cali-bio.com

Sequence

MSWWHHHHH NWNIPPTTQDT TERDCRVSSF
RVKENFDKAR FSGTWYAMAK KDPEGLFLQD
NIVAEFSVDE TGQMSATAKG RVRLLNNWDV

Product Data Sheet

CADMVGTFTD TEDPAETKCYWGADE
GNDDHWIVDT DYDTYAVQYS CRLNLDGTC
ADSYFVFSR DPNGLPPEAQ KIVRQRQEEL
CLARQYRLIV HNGYCDGRSERNLL.

*** For Non-Clinical Research Use Only ***